

ABOUT THIS DOCUMENT

This biogeographic assessment represents the continuation of an ongoing partnership between the National Marine Sanctuary Program (NMSP) and the National Centers for Coastal Ocean Science (NCCOS). The purpose of this collaboration is to provide sanctuary managers with basic information on the distribution of marine flora and fauna relevant to the national marine sanctuaries they manage. This particular work, conducted in collaboration with the Channel Islands National Marine Sanctuary (CINMS) and members of the local research community, builds on a previous assessment developed for California's other three national marine sanctuaries (NOAA, 2003). These efforts were undertaken specifically to support the management plan revision process mandated for each sanctuary. This process evaluates the degree that each sanctuary is meeting its goals and allows an opportunity for the public to determine if there are new directions or issues that they feel the sanctuary should address. One issue raised by the public during the CINMS management plan revision process was whether the sanctuary boundaries should be expanded. A significant portion of this document, therefore, is devoted toward providing a biogeographic assessment of the differing boundary concepts previously developed by CINMS in conjunction with the Sanctuary Advisory Council and general public. This was accomplished by a thorough analysis of the biogeographic datasets provided to the analytical team by the local research community. Additionally, the data gathered, analyses performed, and patterns of distribution observed should provide invaluable information to support science, education, and support other spatially-explicit management decisions.

The results of this assessment are available via both hard copy and CD-ROM. Also available on the CD-ROM are the data utilized to develop the Habitat Suitability Models along with the ArcGIS project files used to develop many of the figures within this report (e.g. species distribution, substrate and oceanographic maps). For more information on this effort please visit the NCCOS Biogeography Team webpage dedicated to this project at: http://biogeo.nos.noaa.gov/projects/assess/ca_nms/cinms/ or direct questions and comments to:

Mark Monaco
Biogeography Team Manager
National Oceanic & Atmospheric Administration
1305 East-West Hwy. (SSMC4, N/SCI-1)
Silver Spring, MD 20910
Phone: (301) 713-3028 x160
Email: mark.monaco@noaa.gov

Or

Chris Mobley
Channel Islands National Marine Sanctuary Manager
National Oceanic & Atmospheric Administration
113 Harbor Way, Suite 150
Santa Barbara, CA 93109
Phone: (805) 884-1465
Email: chris.mobley@noaa.gov

PROJECT TEAM

Satie Airamé-Partnership for Interdisciplinary Studies of Coastal Oceans
M. James Allen-Southern California Coastal Water Research Project Authority
Jay Barlow-Southwest Fisheries Science Center
Ken Buja-National Centers for Coastal Ocean Science
Chris Caldow-National Centers for Coastal Ocean Science
Harry Carter-Carter Biological Consulting
Jenn Caselle-Partnership for Interdisciplinary Studies of Coastal Oceans
John Christensen-National Centers for Coastal Ocean Science
Larry Clafflin-National Centers for Coastal Ocean Science
Randy Clark-National Centers for Coastal Ocean Science
Michael Coyne-National Centers for Coastal Ocean Science
Kate Eschelbach-National Centers for Coastal Ocean Science
Sarah Fangman-Channel Islands National Marine Sanctuary
Glen Ford-R.G. Ford Consulting Company
Karin Forney-Southwest Fisheries Science Center
Steve Gaines-University of California, Santa Barbara
Tracy Gill-National Centers for Coastal Ocean Science
Brian Hatfield-United States Geological Survey
Jamie Higgins-National Centers for Coastal Ocean Science
Olaf Jensen-National Centers for Coastal Ocean Science
Julie Kellner-National Centers for Coastal Ocean Science
Mark Lowry-Southwest Fisheries Science Center
Sarah MacWilliams-Channel Islands National Marine Sanctuary
Chris Mobley-Channel Islands National Marine Sanctuary
Mark Monaco-National Centers for Coastal Ocean Science
Wendy Morrison-Georgia Tech University
Michael Murray-Channel Islands National Marine Sanctuary
Matt Pickett-NOAA Marine and Aviation Operations
Dan Pondella-Occidental College
Lynn Takata-California State Lands Commission
Mitchell Tartt-Office of National Marine Sanctuaries
Jenny Waddell-National Centers for Coastal Ocean Science
Ben Waltenberger-Channel Islands National Marine Sanctuary
Kim Woody-National Centers for Coastal Ocean Science

EXECUTIVE SUMMARY

The priority management goal of the National Marine Sanctuaries Program (NMSP) is to protect marine ecosystems and biodiversity. This goal requires an understanding of broad-scale ecological relationships and linkages between marine resources and physical oceanography to support an ecosystem management approach. The Channel Islands National Marine Sanctuary (CINMS) is currently reviewing its management plan and investigating boundary expansion. A management plan study area (henceforth, Study Area) was described that extends from the current boundary north to the mainland, and extends north to Point Sal and south to Point Dume. Six additional boundary concepts were developed that vary in area and include the majority of the Study Area. The NMSP and CINMS partnered with NOAA's National Centers for Coastal Ocean Science Biogeography Team to conduct a biogeographic assessment to characterize marine resources and oceanographic patterns within and adjacent to the sanctuary. This assessment includes a suite of quantitative spatial and statistical analyses that characterize biological and oceanographic patterns in the marine region from Point Sal to the U.S.-Mexico border. These data were analyzed using an index which evaluates an ecological "cost-benefit" within the proposed boundary concepts and the Study Area.

The sanctuary resides in a dynamic setting where two oceanographic regimes meet. Cold northern waters mix with warm southern waters around the Channel Islands creating an area of transition that strongly influences the regions oceanography. In turn, these processes drive the biological distributions within the region. This assessment analyzes bathymetry, benthic substrate, bathymetric life-zones, sea surface temperature, primary production, currents, submerged aquatic vegetation, and kelp in the context of broad-scale patterns and relative to the proposed boundary concepts and the Study Area. Boundary cost-benefit results for these parameters were variable due to their dynamic nature; however, when analyzed in composite the Study Area and Boundary Concept 2 were considered the most favorable.

Biological data were collected from numerous resource agencies and university scientists for this assessment. Fish and invertebrate trawl data were used to characterize community structure. Habitat suitability models were developed for 15 species of macroinvertebrates and 11 species of fish that have significant ecological, commercial, or recreational importance in the region and general patterns of ichthyoplankton distribution are described. Six surveys of ship and plane at-sea surveys were used to model marine bird diversity from Point Arena to the U.S.-Mexico border. Additional surveys were utilized to estimate density and colony counts for nine bird species. Critical habitat for western snowy plover and the location of California least tern breeding pairs were also analyzed. At-sea surveys were also used to describe the distribution of 14 species of cetaceans and five species of pinnipeds. Boundary concept cost-benefit indices revealed that Boundary Concept 2 and the Study Area were most favorable for the majority of the species-specific analyses. Boundary Concept 3 was most favorable for bird diversity across the region. Inadequate spatial resolution for fish and invertebrate community data and incompatible sampling effort information for bird and mammal data precluded boundary cost-benefit analysis.

The final chapter integrates data and analyses from each of the preceding chapters utilizing two separate approaches. Cost-benefit indices were ranked for each biological group and for the oceanographic/physical parameters to provide a consistent and comprehensive evaluation of the boundary concepts. The Study Area and Boundary Concept 2 (see Chapter 1) ranked highest for the bird, fish, and mammal groups, as well as all the data in composite. The Study Area also ranked highest for macroinvertebrates. Second, select spatial data were integrated, based on data compatibility and spatial range, to identify areas of spatial coincidence which may reflect ecosystem "hotspots". Habitat suitability models for fish and macroinvertebrates, along with bird and mammal sightings information were utilized to evaluate this spatial coincidence. Areas of highest spatial coincidence most closely resemble the spatial delineation for the Study Area and also include a broad area from the mainland south through San Clemente Island.

Integration results highlight the Channel Islands and the area extending north to the mainland to Point Conception as an important ecosystem that supports a diverse array of biological communities. The boundary concepts that were favorably ranked incorporated large areas of the coastal mainland, due in part to the nearshore affinity exhibited by many of the analyzed species. Deep offshore environments away from the Channel Islands were

correspondingly less favorable. Both the Study Area and Boundary Concept 2 are characterized by areas of increased upwelling, dynamic surface currents and eddies, and persistent thermal fronts. These concepts also include large areas of important habitats such as kelp, seagrasses, and wetlands along with a mixture of deep and shallow waters that many species depend on for all or part of their life cycles.

In compliance with the National Environmental Policy Act, the National Marine Sanctuary Program will incorporate this assessment with cultural and socio-economic analyses to prepare a Supplemental Environmental Impact Statement to fully analyze boundary change concepts.

TABLE OF CONTENTS

CHAPTER 1 INTRODUCTION	1
1.1 PROJECT BACKGROUND	1
1.2 INTRODUCTION TO BIOGEOGRAPHY	5
1.3 BIOGEOGRAPHY OF THE WEST COAST	6
Marine Benthic Invertebrates	7
Marine Fishes	8
Seabirds and Shorebirds	9
Marine Mammals	10
1.4 THE FOUR-STEP ASSESSMENT PROCESS	11
Species Selection	11
Data Collection and Synthesis	11
Metric Development	13
Analyses Review	13
1.5 ASSESSMENT OUTLINE	14
LITERATURE CITED	15
CHAPTER 2 PHYSICAL AND OCEANOGRAPHIC SETTING	17
2.1 PHYSICAL ENVIRONMENT AND GEOLOGY	17
2.2 CLIMATE AND METEOROLOGY	18
2.3 PHYSICAL OCEANOGRAPHY	18
Offshore Ocean Currents	19
Waves	21
Long-Term Climate Perturbations	21
2.4 PHYSIOGRAPHIC COMPLEXITY	21
Data and Methods	21
Broad-scale Patterns	22
Analysis of Boundary Concepts	23
Summary	24
2.5 BENTHIC SUBSTRATE	26
Data and Methods	26
Broad-scale Patterns	27
Analysis of Boundary Concepts	28
Summary	30
2.6 BATHYMETRIC LIFE-ZONES	31
Data and Methods	31
Broad-scale Patterns	31
Analysis of Boundary Concepts	32
Summary	33
2.7 SEA SURFACE TEMPERATURE AND FRONTAL BOUNDARIES	33
Data and Methods	33
Broad-scale Patterns	34
Analysis of Boundary Concepts	35
Summary	37
2.8 SURFACE CHLOROPHYLL AND OCEAN CURRENTS	38
Data and Methods	38
Broad-scale Patterns	40
Analysis of Boundary Concepts	42
Summary	45
2.9 SUBMERGED AQUATIC VEGETATION: EELGRASS AND SURFGRASS	47
Data and Methods	47
Broad-scale Patterns	48
Analysis of Boundary Concepts	50
Summary	50

2.10 KELP DISTRIBUTION	50
Data and Methods	50
Broad-scale Patterns	51
Analysis of Boundary Concepts	52
Summary	52
LITERATURE CITED	52
CHAPTER 3 BIOGEOGRAPHY OF MARINE MACROINVERTEBRATES	57
3.1 SINGLE SPECIES HABITAT SUITABILITY MODELS (HSM)	57
Data and Methods	57
Rock crabs (<i>Cancer</i> spp.)	58
Black abalone (<i>Haliotis cracherodii</i>)	60
Red abalone (<i>Haliotis rufescens</i>)	61
White abalone (<i>Haliotis sorenseni</i>)	63
California market squid (<i>Loligo opalescens</i>)	65
Sheep crab (<i>Loxorhynchus grandis</i>)	67
Spot shrimp (<i>Pandalus platyceros</i>)	68
Ridgeback rock shrimp (<i>Sicyonia ingentis</i>)	71
California spiny lobster (<i>Panulirus interruptus</i>)	72
California sea cucumber (<i>Parastichopus californicus</i>)	73
Warty sea cucumber (<i>Parastichopus parvimensis</i>)	75
Red sea urchin (<i>Strongylocentrotus franciscanus</i>)	76
Purple sea urchin (<i>Strongylocentrotus purpuratus</i>)	78
3.2 MACROINVERTEBRATE ASSEMBLAGE STRUCTURE	80
Data and Methods	80
Broad-Scale Patterns	81
Analysis of Boundary Concepts	82
Summary	85
LITERATURE CITED	85
CHAPTER 4 BIOGEOGRAPHY OF MARINE FISHES	89
4.1 SINGLE SPECIES HABITAT SUITABILITY MODELS (HSM)	89
Data and Methods	89
Thresher shark (<i>Alopias vulpinus</i>)	90
Tope (<i>Galeorhinus galeus</i>)	92
Leopard shark (<i>Triakis semifasciata</i>)	95
Pacific angel shark (<i>Squatina californica</i>)	97
Pacific sardine (<i>Sardinops sagax</i>)	98
Northern anchovy (<i>Engraulis mordax</i>)	98
Bocaccio (<i>Sebastes paucispinis</i>)	99
Cowcod (<i>Sebastes levis</i>)	101
Lingcod (<i>Ophiodon elongatus</i>)	104
Giant seabass (<i>Stereolipis gigas</i>)	107
California sheephead (<i>Semicossyphus pulcher</i>)	108
California halibut (<i>Paralichthys californicus</i>)	110
4.2 FISH ASSEMBLAGE STRUCTURE	112
Data and Methods	112
Broad-scale Patterns	115
Diversity	115
Assemblages	116
Analysis of Boundary Concepts	125
Summary	125
4.3 ICHTHYOPLANKTON	125
Data and Methods	125
Broad-scale Patterns	128
California halibut (<i>Paralichthys californicus</i>)	128
Bocaccio (<i>Sebastes paucispinis</i>)	128

Pacific sardine (<i>Sardinops sagax</i>).....	129
Northern anchovy (<i>Engraulis mordax</i>).....	129
Analysis of Boundary Concepts	130
Summary	130
LITERATURE CITED	131
CHAPTER 5 BIOGEOGRAPHY OF MARINE BIRDS	135
5.1 MARINE BIRD SINGLE SPECIES ANALYSIS	135
Data and Methods	135
Computer Database Analysis System (CDAS)	135
Xantus's Murrelet Telemetry Study	138
Broad-scale Patterns and Analysis of Boundary Concepts	138
Ashy storm-petrel (<i>Oceanodroma homochroa</i>)	138
California brown pelican (<i>Pelecanus occidentalis californicus</i>)	140
Double-crested cormorant (<i>Phalacrocorax auritus</i>)	142
Brandt's cormorant (<i>Phalacrocorax penicillatus</i>)	142
Pelagic cormorant (<i>Phalacrocorax pelagicus</i>)	145
Western snowy plover (<i>Charadrius alexandrinus nivosus</i>)	147
Black oystercatcher (<i>Haematopus bachmani</i>)	148
California least tern (<i>Sterna antillarum browni</i>)	149
Pigeon guillemot (<i>Cepphus columba</i>)	149
Xantus's murrelet (<i>Synthliboramphus hypoleucus</i>)	150
Cassin's auklet (<i>Ptychoramphus aleuticus</i>)	153
Other Marine Birds	156
Summary	156
5.2 MARINE BIRD DIVERSITY	157
Data and Methods	157
Broad-scale Patterns	158
Analysis of Boundary Concepts	160
Summary	162
LITERATURE CITED	163
CHAPTER 6 BIOGEOGRAPHY OF MARINE MAMMALS	167
6.1 CETACEAN SINGLE SPECIES ANALYSIS	167
Data and Methods	167
SWFSC Shipboard Suveys	168
SWFSC Aerial Surveys	171
Computer Database Analysis System (CDAS)	171
SWFSC Bottlenose Dolphin Aerial Survey	171
Broad-scale Patterns and Analysis of Boundary Concepts	172
Blue whale (<i>Balaenoptera musculus</i>)	172
Bottlenose dolphin (<i>Tursiops truncatus</i>)	173
Long-beaked and short-beaked common dolphins (<i>Delphinus</i> spp.)	175
Gray whale (<i>Eschrichtius robustus</i>)	178
Humpback whale (<i>Megaptera novaeangliae</i>)	178
Killer whale (<i>Orcinus orca</i>)	179
Pacific white-sided dolphin (<i>Lagenorhynchus obliquidens</i>)	180
Risso's dolphin (<i>Grampus griseus</i>)	181
Additional Cetaceans	183
Dall's porpoise (<i>Phocoenoides dalli</i>)	183
Fin whale (<i>Balaenoptera physalus</i>)	183
Harbor porpoise (<i>Phocoena phocoena</i>)	184
Minke whale (<i>Balaenoptera acutorostrata</i>)	185
Northern righ-whale dolphin (<i>Lissodelphis borealis</i>)	185
Summary	185
6.2 PINNIPEDS AND SOUTHERN SEA OTTER	186
Pinnipedia (seals, sea lions, and fur seals)	186
Carnivora	187
Data and Methods	187

California sea lion (<i>Zalophus californianus californianus</i>)	187
Pacific harbor seal (<i>Phoca vitulina richardsi</i>)	188
Northern elephant seal (<i>Mirounga angustirostris</i>)	188
Northern fur seal (<i>Callorhinus ursinus</i>).....	188
Southern sea otter (<i>Enhydra lutris nereis</i>).....	190
Broad-scale Patterns and Analysis of Boundary Concepts	188
California sea lion.....	188
Pacific harbor seal.....	190
Northern elephant seal	191
Southern sea otter.....	193
Summary	194
LITERATURE CITED	196
CHAPTER 7 INTEGRATION OF ANALYSES	199
7.1 NUMERICAL INTEGRATION	199
Data and Methods	199
Analysis of Boundary Concepts	199
7.2 SPATIAL INTEGRATION	201
Data and Methods	201
Broad-scale Patterns and Analysis of Boundary Concepts	202
Summary	205
7.3 THREATENED AND ENDANGERED SPECIES	206
White abalone	206
Western snowy plover	206
California brown pelican	206
California least tern	207
Southern sea otter.....	207
Blue whale.....	207
Humpback whale.....	208
Summary	208
LITERATURE CITED	209
APPENDIX A	211
ACKNOWLEDGEMENTS	215

List of Tables

Table 1.1.1. Total area, perimeter, and amount of mainland coastline included for the NAC, Study Area, and six boundary concepts.....	4
Table 2.4.1. Analysis of boundary concepts.	26
Table 2.5.1. Area (km ²) and percentage of total area for habitat types within boundary concepts.	28
Table 2.5.2. Analysis of habitat diversity within boundary concepts.	30
Table 2.6.1. Analysis of life-zone evenness within boundary concepts.	33
Table 2.8.1. Analysis of chlorophyll within boundary concepts.	45
Table 2.9.1. Analysis of SAV distribution within boundary concepts.	50
Table 2.10.1. Analysis of kelp distribution within boundary concepts.	52
Table 3.1.1. Invertebrate species of interest for the CINMS biogeographic assessment.	57
Table 3.1.2. Analysis of rock crab habitat suitability within boundary concepts.	60
Table 3.1.3. Analysis of black abalone habitat suitability within boundary concepts.	62
Table 3.1.4. Analysis of red abalone habitat suitability within boundary concepts.	63
Table 3.1.5. Analysis of white abalone habitat suitability within boundary concepts.	65
Table 3.1.6. Analysis of California market squid habitat suitability within boundary concepts.	67
Table 3.1.7. Analysis of sheep crab habitat suitability within boundary concepts.	68
Table 3.1.8. Analysis of spot shrimp habitat suitability within boundary concepts.	70
Table 3.1.9. Analysis of ridgeback shrimp habitat suitability within boundary concepts.	72
Table 3.1.10. Analysis of California spiny lobster habitat suitability within boundary concepts.	74
Table 3.1.11. Analysis of California sea cucumber habitat suitability within boundary concepts.	75
Table 3.1.12. Analysis of warty sea cucumber habitat suitability within boundary concepts.	77
Table 3.1.13. Analysis of red sea urchin habitat suitability within boundary concepts.	78
Table 3.1.14. Analysis of purple sea urchin habitat suitability within boundary concepts.	79
Table 3.2.1. Species assemblage results for SCCWRP survey data (1994, 1998) using the Bray-Curtis dissimilarity metric with average means clustering.	82
Table 3.2.2. Mean frequency of occurrence for each SCCWRP site group.	83
Table 4.1.1. Fish species of interest for the CINMS biogeographic assessment.	89
Table 4.1.2. Analysis of adult thresher shark habitat suitability within boundary concepts.	92
Table 4.1.3. Analysis of juvenile thresher shark habitat suitability within boundary concepts.	92
Table 4.1.4. Analysis of adult tope habitat suitability within boundary concepts.	94
Table 4.1.5. Analysis of juvenile tope habitat suitability within boundary concepts.	95
Table 4.1.6. Analysis of adult leopard shark habitat suitability within boundary concepts.	96
Table 4.1.7. Analysis of Pacific angel shark habitat suitability within boundary concepts.	98
Table 4.1.8. Analysis of adult bocaccio habitat suitability within boundary concepts.	101
Table 4.1.9. Analysis of juvenile bocaccio habitat suitability within boundary concepts.	101
Table 4.1.10. Analysis of adult cowcod habitat suitability within boundary concepts.	103
Table 4.1.11. Analysis of juvenile cowcod habitat suitability within boundary concepts.	104
Table 4.1.12. Analysis of adult lingcod habitat suitability within boundary concepts.	106
Table 4.1.13. Analysis of juvenile lingcod habitat suitability within boundary concepts.	106
Table 4.1.14. Analysis of giant seabass habitat suitability within boundary concepts.	108
Table 4.1.15. Analysis of California sheephead habitat suitability within boundary concepts.	110
Table 4.1.16. Analysis of California halibut habitat suitability within boundary concepts.	112
Table 4.2.1. Summary of the datasets used to assess fish diversity and species assemblages.	113
Table 4.2.2. Species assemblage results for the RecFIN CPFV data using the Bray-Curtis dissimilarity metric with average means clustering.	117
Table 4.2.3. Mean frequency of occurrence for each recreational site group.	119
Table 4.2.4. Species assemblage results for the SCCWRP trawl data using the Bray-Curtis dissimilarity metric with average means clustering.	120
Table 4.2.5. Mean frequency of occurrence for SCCWRP site groups.	121
Table 4.2.6. Species assemblage results for the NMFS GSP data using Bray-Curtis dissimilarity metric with average means clustering.	122
Table 4.2.7. Mean frequency of occurrence for NMFS GSP site groups.	123

Table 4.2.8. Species assemblage results for kelp visual census surveys using Bray-Curtis dissimilarity metric with average means clustering	123
Table 4.2.9. Mean frequency of occurrence for kelp visual census site groups	124
Table 5.0.1. Status, habitat locations, and seasonal use of the requested marine birds in the Study Area.....	136
Table 5.1.1. Data used for the analysis of marine bird distributions presented in this chapter.....	137
Table 5.1.2. Ashy storm-petrel. At-sea sightings, effort, density, and Optimal-Area Index (OAI) for each boundary concept.....	140
Table 5.1.3. California brown pelican. At-sea sightings, effort, density, and Optimal-Area Index (OAI) for each boundary concept.....	142
Table 5.1.4. Double-crested cormorant. At-sea sightings, effort, density, and Optimal-Area Index (OAI) for each boundary concept.....	144
Table 5.1.5. Brandt's cormorant. At-sea sightings, effort, density, and Optimal-Area Index (OAI) for each boundary concept.....	145
Table 5.1.6. Pelagic cormorant. At-sea sightings, effort, density, and Optimal-Area Index (OAI) for each boundary concept.....	147
Table 5.1.7. Pelagic cormorant. Colony counts (total individuals) and Optimal-Area Index (OAI) for each boundary concept.....	147
Table 5.1.8. Western snowy plover. Critical habitat for the Pacific coast population of western snowy plover.....	148
Table 5.1.9. Black oystercatcher. Colony counts (total individuals) and Optimal-Area Index (OAI) for each boundary concept	148
Table 5.1.10. California least tern. The maximum number of breeding pairs observed at each nesting site surveyed from 2001-2003 for each boundary concept (CDFG, DRAFT data)..	149
Table 5.1.11. Pigeon guillemot. At-sea sightings, effort, density, and Optimal-Area Index (OAI) for each boundary concept	150
Table 5.1.12. Pigeon guillemot. Colony counts (total individuals) and Optimal-Area Index (OAI) for each boundary concept.....	151
Table 5.1.13. Xantus's murrelet. At-sea sightings, effort, density, and Optimal-Area Index (OAI) for each boundary concept.....	152
Table 5.1.14. Xantus's murrelet. Telemetry relocations and Optimal-Area Index (OAI) for each boundary concept.....	153
Table 5.1.15. Cassin's auklet. At-sea sightings, effort, density, and Optimal-Area Index (OAI) for each boundary concept.....	156
Table 5.2.1. Summary of the six surveys that were used in the analysis of marine bird diversity.....	157
Table 5.2.2. Analysis of bird diversity within boundary concepts	162
Table 6.1.1. Summary of marine mammal field surveys examined in this chapter.....	168
Table 6.1.2. Line transect parameters used to estimate the abundance of selected cetaceans within the six proposed boundary concepts, the current CINMS boundary, and the Study Area.....	168
Table 6.1.3. Blue whale. Sightings, estimated density and abundance, coefficient of variation (CV) and upper and lower 95% confidence limits for the abundance estimate, and the Optimal Area Index (OAI) for the six proposed boundary concepts, the No Action Concept (NAC), and the Study Area (SA)	173
Table 6.1.4. Coastal bottlenose dolphin. Sightings, mean encounter rate, and estimated abundance for four proposed boundary concepts and the Study Area.....	175
Table 6.1.5. Long-beaked common dolphin. Sightings, estimated density and abundance, coefficient of variation (CV) and upper and lower 95% confidence limits for the abundance estimate (corrected for unidentified common dolphin sightings), and the Optimal Area Index (OAI) for the six proposed boundary concepts, the No Action Concept (NAC), and the Study Area (SA)	177

Table 6.1.6. Short-beaked common dolphin. Sightings, estimated density and abundance, coefficient of variation (CV) and upper and lower 95% confidence limits for the abundance estimate (corrected for unidentified common dolphin sightings), and the Optimal Area Index (OAI) for the six proposed boundary concepts, the No Action Concept (NAC), and the Study Area (SA).	177
Table 6.1.7. Humpback whale. Sightings, estimated density and abundance, coefficient of variation (CV) and upper and lower 95% confidence limits for the abundance estimate for the six proposed boundary concepts, the No Action Concept (NAC), and the Study Area (SA).	179
Table 6.1.8. Risso's dolphin. Sightings, estimated density and abundance, coefficient of variation (CV) and upper and lower 95% confidence limits for the abundance estimate, and the Optimal Area Index (OAI) for the six proposed boundary concepts, the No Action Concept (NAC), and the Study Area (SA)	182
Table 6.2.1. Summary of pinniped and sea otter surveys used in this chapter	187
Table 6.2.2. Pacific Harbor seal. Total number of individuals, total area, and OAI for each boundary concept for SWFSC and CDFG 2002 surveys	192
Table 6.2.3. Southern Sea otter. Total number of individuals, mainland encounter rates, total area, and OAI for each boundary concept for Fall 2001 and Spring 2002 surveys.....	195
Table 7.1.1. Absolute OAI rankings for individual, groups, and the composite for all analyses	200
Table 7.1.2. Results of Kruskal-Wallis ranks sums test.	201
Table 7.3.1. Analysis of white abalone habitat suitability within boundary concepts	207
Table 7.3.2. Critical habitat for the Pacific coast population of western snowy plover.....	206
Table 7.3.3. California brown Pelican sightings and density OAI	206
Table 7.3.4. California least tern breeding pairs observed within boundary concepts, 2001-2003.....	207
Table 7.3.5. Spring and fall abundance estimates and OAI for southern sea otter.....	207
Table 7.3.6. Sightings, estimated abundance and density, and OAI for blue whales	207
Table 7.3.7. Sightings, estimated abundance and density for humpback whales	208
Table 7.3.8. Ranked OAI for the federally listed threatened or endangered species	208

List of Figures

Figure 1.1.1. Map of the Channel Islands and specific coastal locations in the surrounding region of interest	1
Figure 1.1.2a. Spatial delineations for the Study Area and boundary Concepts 1-4.....	3
Figure 1.1.2b. Spatial delineation for boundary Concept 5	4
Figure 1.3.1. Pelagic life-zones off southern California	7
Figure 1.3.2. Benthic life-zones off southern California.....	8
Figure 1.3.3. Latitudinal range endpoints for 539 species of marine benthic invertebrates	9
Figure 1.3.4. Latitudinal range endpoints for 294 species of marine fishes	10
Figure 1.3.5. Latitudinal range endpoints for 132 shore and seabird species.....	11
Figure 1.3.6. Latitudinal range endpoints for 49 marine mammal species.....	12
Figure 1.4.1. A hypothetical set of three boundary concepts and the ecological value of the area contained within them	14
Figure 1.4.2. Trend in values of absolute and relative metrics and the OAI for the hypothetical example shown in Figure 1.1.9.....	14
Figure 2.1.1. Bathymetric and geologic features for the region of interest.....	17
Figure 2.4.1. Map of physiographic complexity along the California coast (ranging from 32° to 39° north latitude)	24
Figure 2.4.2. Physiographic complexity within boundary concepts	25
Figure 2.4.3. Linear regression functions between concept area (km ²) and mean physiographic complexity (left) and and the area (km ²) of high physiographic complexity (right).....	26
Figure 2.5.1. Benthic habitat types for the marine waters off California	27
Figure 2.5.2. Location of hard substrate from NMFS and MMS datasets	29
Figure 2.5.3. Benthic habitat types for the marine waters off southern California	29

Figure 2.5.4. Regression of habitat diversity and total area for the current and proposed boundary concepts.....	30
Figure 2.5.5. Evenness and richness estimates for habitat types within boundary concepts	30
Figure 2.6.1. Distribution of benthic substrate types (hard or soft) off southern California	31
Figure 2.6.2. Distribution of bathymetric zones (m) off southern California	32
Figure 2.6.3. Regression of habitat evenness and total area for the current and proposed boundary concepts.....	33
Figure 2.7.1. Seasonal mean sea surface temperature (January, April, July, and October) averaged across years from 1993 to 2003	35
Figure 2.7.2. Sea surface temperature variance, calculated from mean monthly values for the period January 1993 to May 2003	36
Figure 2.7.3. Left panel shows an example interpolated SST grid from GMT for January 1993.....	36
Figure 2.7.4. Persistence of SST fronts through time	37
Figure 2.7.5. Areas with greatest SST front activity through time	37
Figure 2.7.6. Linear regression function between concept area (km ²) and mean persistence of SST fronts	38
Figure 2.7.7. Linear regression function between total concept area (km ²) and area of high SST frontal persistence.....	38
Figure 2.8.1. Average [ChlA] for grid cells in which data was present during November 1997-December 1999	39
Figure 2.8.2. Interpolated mean chlorophyll concentration [ChlA] for the period between November 1997 and December 1999. Model extent ranges from 28°- 49° latitude.....	40
Figure 2.8.3. Interpolated mean chlorophyll concentration [ChlA] for the period between November 1997 and December 1999.....	41
Figure 2.8.4. Monthly mean [ChlA] for 1999. Estimates derived from data contained within the Study Area boundary.....	42
Figures 2.8.5a. Mean chlorophyll concentration [ChlA] for the months January to June 1999. Altimetry derived surface current velocity vectors for the same time periods are superimposed on the [ChlA] surface.....	43
Figures 2.8.5b. Mean chlorophyll concentration [ChlA] for the months of July to December 1999 (left to right). Altimetry derived surface current velocity vectors for the same time periods are superimposed on the [ChlA] surface.....	44
Figures 2.8.6. Interpolated mean chlorophyll concentration [ChlA] for the period between November 1997 and December 1999 with boundary concepts overlain	46
Figure 2.8.7. Linear regression function between concept area (km ²) and mean [ChlA]	47
Figure 2.8.8. Linear regression function between the total concept area (km ²) and area of high [ChlA]	47
Figure 2.9.1. Distribution of submerged aquatic vegetation within coastal California waters.....	48
Figure 2.9.2. Distribution of submerged aquatic vegetation within the proposed boundary concepts in southern California.....	49
Figure 2.9.3. Linear regression function between area of SAV distribution and total area of boundary concepts.....	50
Figure 2.10.1. Kelp distribution based on aerial and multispectral surveys conducted by CDFG during 1989, 1999, and 2002. Kelp polygons have been enlarged for better viewing	51
Figure 2.10.2. Kelp distribution off southern California	51
Figure 2.10.3. Regression of kelp area and total area for the current and proposed boundary concepts.....	52
Figure 3.1.1. CDFG commercial 10x10 nm fishery landings grids. Mean monthly landings (pounds) were reported within designated grids from 1996-2002.....	58
Figure 3.1.2. Rock crab (<i>Cancer</i> spp.) habitat suitability off central and southern California.....	59
Figure 3.1.3. Rock crab (<i>Cancer</i> spp.) habitat suitability off southern California	59
Figure 3.1.4. Regression of highly suitable habitat area for rock crabs and total area for the current and proposed boundary concepts	60

Figure 3.1.5. Black abalone habitat suitability off southern California.....	61
Figure 3.1.6. Black abalone commercial landings data from CDFG CMASTR database, 1990-1993, superimposed over predicted habitat suitability.....	61
Figure 3.1.7. Regression of highly suitable habitat area for black abalone and total area for the current and proposed boundary concepts	61
Figure 3.1.8. Red abalone habitat suitability off southern California	62
Figure 3.1.9. Red abalone commercial landings data from CDFG CMASTR database, 1990-1999, superimposed over predicted habitat suitability	63
Figure 3.1.10. Regression of highly suitable habitat area for red abalone and total area for the current and proposed boundary concepts	63
Figure 3.1.11. White abalone habitat suitability off southern California.....	64
Figure 3.1.12. White abalone commercial landings data from CDFG CMASTR database, 1984-1999, superimposed over predicted habitat suitability.....	64
Figure 3.1.13. Regression of highly suitable habitat area for white abalone and total area for the current and proposed boundary concepts	65
Figure 3.1.14. California market squid habitat suitability off central and southern California.....	66
Figure 3.1.15. Location of NMFS trawls (1999-2002) and squid mean log abundance superimposed over predicted habitat suitability	66
Figure 3.1.16. California market squid habitat suitability off southern California.....	66
Figure 3.1.17. Regression of highly suitable habitat area for California market squid and total area for the current and proposed boundary concepts	67
Figure 3.1.18. Sheep crab habitat suitability off central and southern California	67
Figure 3.1.19. Sheep crab commercial landings data from CDFG CMASTR database, 1996-2000, superimposed over predicted habitat suitability.....	68
Figure 3.1.20. Sheep crab habitat suitability off southern California	68
Figure 3.1.21. Regression of highly suitable habitat area for sheep crab and total area for the current and proposed boundary concepts	68
Figure 3.1.22. Spot shrimp habitat suitability off central and southern California	69
Figure 3.1.23. Spot shrimp commercial landings data from CDFG Commercial Trap Logs, 1994-2001, superimposed over predicted habitat suitability.....	69
Figure 3.1.24. Spot shrimp commercial landings data from CDFG Commercial Trawl Logs, 1994-2001, superimposed over predicted habitat suitability.....	70
Figure 3.1.25. Spot shrimp habitat suitability off southern California	70
Figure 3.1.26. Regression of highly suitable habitat area for spot shrimp and total area for the current and proposed boundary concepts	70
Figure 3.1.27. Ridgeback shrimp habitat suitability off central and southern California.....	71
Figure 3.1.28. Location of SCCWRP trawls (1994, 1998) and ridgeback shrimp mean abundance superimposed over predicted habitat suitability	71
Figure 3.1.29. Ridgeback shrimp habitat suitability off southern California.....	71
Figure 3.1.30. Regression of highly suitable habitat area for ridgeback shrimp and total area for the current and proposed boundary concepts	72
Figure 3.1.31. California spiny lobster habitat suitability off central and southern California	73
Figure 3.1.32. California spiny lobster commercial landings data from CDFG Commercial Logs, 1998-2002, superimposed over predicted habitat suitability.....	73
Figure 3.1.33. California spiny lobster habitat suitability off southern California	73
Figure 3.1.34. Regression of highly suitable habitat area for California spiny lobster and total area for the current and proposed boundary concepts	74
Figure 3.1.35. California sea cucumber habitat suitability off central and southern California.....	74
Figure 3.1.36. Location of SCCWRP trawls (1994, 1998) and California sea cucumber mean abundance superimposed over predicted habitat suitability	74
Figure 3.1.37. California sea cucumber habitat suitability off southern California.....	75
Figure 3.1.38. Regression of highly suitable habitat area for California sea cucumber and total area for the current and proposed boundary concepts	75
Figure 3.1.39. Warty sea cucumber habitat suitability off central and southern California.....	76

Figure 3.1.40. Warty sea cucumber habitat suitability off southern California.....	76
Figure 3.1.41. Regression of highly suitable habitat area for warty sea cucumber and total area for the current and proposed boundary concepts	76
Figure 3.1.42. Red sea urchin habitat suitability off central and southern California	77
Figure 3.1.43. Red sea urchin habitat suitability off southern California	77
Figure 3.1.44. Regression of highly suitable habitat area for red sea urchin and total area for the current and proposed boundary concepts	78
Figure 3.1.45. Purple sea urchin habitat suitability off central and southern California.....	78
Figure 3.1.46. Purple sea urchin habitat suitability off southern California	79
Figure 3.1.47. Regression of highly suitable habitat area for purple sea urchin and total area for the current and proposed boundary concepts	79
Figure 3.2.1. Invertebrate diversity for individual SCCWRP trawls during 1994 and 1998	81
Figure 3.2.2. Location of site groups for SCCWRP survey data (1994, 1998).....	81
Figure 3.2.3. Overlay of invertebrate diversity and CINMS boundary concepts.....	84
Figure 4.1.1. Adult thresher shark habitat suitability off central and southern California.....	91
Figure 4.1.2. Juvenile thresher shark habitat suitability off central and southern California	91
Figure 4.1.3. Adult thresher shark habitat suitability off southern California	91
Figure 4.1.4. Regression of total habitat area for adult thresher shark and total area within the current and proposed boundary concepts	91
Figure 4.1.5. Juvenile thresher shark habitat suitability off southern California	92
Figure 4.1.6. Regression of total habitat area for juvenile thresher shark and total area within the current and proposed boundary concepts	92
Figure 4.1.7. Adult tope habitat suitability probability (HSP) off central and southern California	93
Figure 4.1.8. Juvenile tope habitat suitability probability (HSP) off central and southern California	93
Figure 4.1.9. Adult tope habitat suitability probability (HSP) off southern California	93
Figure 4.1.10. Regression of high probability habitat area for adult tope and total area within the current and proposed boundary concepts	94
Figure 4.1.11. Juvenile tope habitat suitability probability (HSP) off southern California	94
Figure 4.1.12. Regression of total habitat area for juvenile tope and total area within the current and proposed boundary concepts	94
Figure 4.1.13. Adult leopard shark habitat suitability probability (HSP) off central and southern California.....	95
Figure 4.1.14. Adult leopard shark habitat suitability probability (HSP) off southern California	96
Figure 4.1.15. Regression of high probability habitat area for adult leopard shark and total area within the current and proposed boundary concepts	96
Figure 4.1.16. Pacific angel shark habitat suitability off central and southern California	97
Figure 4.1.17. Pacific angel shark habitat suitability off southern California	97
Figure 4.1.18. Regression of highly suitable habitat area for Pacific angel shark and total area within the current and proposed boundary concepts	97
Figure 4.1.19. Adult bocaccio habitat suitability probability (HSP) off central and southern California.....	99
Figure 4.1.20. Juvenile bocaccio habitat suitability probability (HSP) off central and southern California.....	99
Figure 4.1.21. Adult bocaccio habitat suitability probability (HSP) off southern California	100
Figure 4.1.22. Regression of high probability habitat area for adult bocaccio and total area within the current and proposed boundary concepts	100
Figure 4.1.23. Juvenile bocaccio habitat suitability probability (HSP) off southern California	100
Figure 4.1.24. Regression of high probability habitat area for juvenile bocaccio and total area within the current and proposed boundary concepts	100
Figure 4.1.25. Adult cowcod habitat suitability probability (HSP) off central and southern California.....	102
Figure 4.1.26. Juvenile cowcod habitat suitability probability (HSP) off central and southern California.....	102
Figure 4.1.27. Adult cowcod habitat suitability probability (HSP) off southern California	102

Figure 4.1.28. Regression of high probability habitat area for adult cowcod and total area within the current and proposed boundary concepts	102
Figure 4.1.29. Juvenile cowcod habitat suitability probability (HSP) off central and southern California	103
Figure 4.1.30. Regression of high probability habitat area for juvenile cowcod and total area within the current and proposed boundary concepts	103
Figure 4.1.31. Adult lingcod habitat suitability probability (HSP) off central and southern California	105
Figure 4.1.32. Juvenile lingcod habitat suitability probability (HSP) off central and southern California	105
Figure 4.1.33. Adult lingcod habitat suitability probability (HSP) in nearshore and offshore waters of southern California	105
Figure 4.1.34. Regression of high probability habitat area for adult lingcod and total area within the current and proposed boundary concepts	105
Figure 4.1.35. Juvenile lingcod habitat suitability probability (HSP) in nearshore and offshore waters of southern California	106
Figure 4.1.36. Regression of high probability habitat area for juvenile lingcod and total area within the current and proposed boundary concepts	106
Figure 4.1.37. Giant seabass habitat suitability off central and southern California	107
Figure 4.1.38. Giant seabass habitat suitability off southern California	107
Figure 4.1.39. Regression of highly suitable habitat area for giant seabass and total area within the current and proposed boundary concepts	108
Figure 4.1.40. California sheephead habitat suitability off central and southern California	109
Figure 4.1.41. California sheephead landings data from CDFG's Commercial Passenger Fishing Vessel (CPFV) database, 1998-2002, superimposed over predicted habitat suitability	109
Figure 4.1.42. California sheephead habitat suitability off southern California	109
Figure 4.1.43. Regression of highly suitable habitat area for California sheephead and total area within the current and proposed boundary concepts	112
Figure 4.1.44. California halibut habitat suitability off central and southern California	111
Figure 4.1.45. Abundance of California halibut captured in SCCWRP trawls superimposed over predicted habitat suitability	111
Figure 4.1.46. California halibut landings data from CDFG's Commercial Passenger Fishing Vessel (CPFV) database, 1998-2000, superimposed over predicted habitat suitability	111
Figure 4.1.47. California halibut habitat suitability off southern California	111
Figure 4.1.48. Regression of highly suitable habitat area for California halibut and total area within the current and proposed boundary concepts	112
Figure 4.2.1. Fish diversity calculated for individual RecFin hook and line trips (left) and mean diversity for trips within 5x5 minute grids (right)	115
Figure 4.2.2. Fish diversity calculated for individual SCCWRP trawls (left) and mean diversity of trawls within 5x5 minute grids (right)	116
Figure 4.2.3. Fish diversity calculated for individual NMFS GSP trawls (left) and mean diversity of trawls within 5x5 minute grids (right)	116
Figure 4.2.4. Fish diversity calculated for individual kelp visual census surveys (left) and mean diversity of surveys within 5x5 minute grids (right)	117
Figure 4.2.5. Composite fish diversity (mean of standardized values across the four datasets (left) and effort within 5x5 minute grids (right)	117
Figure 4.2.6. Location of site groups, RecFIN CPFV data	118
Figure 4.2.7. Location of site groups for the RecFIN CPFV data within southern California (left) and around Anacapa Island.	119
Figure 4.2.8. Location of site groups, SCCWRP.	121
Figure 4.2.9. Location of site groups, NMFS GSP.	121
Figure 4.2.10. Location of site groups, kelp visual census surveys.	121
Figure 4.2.11. Composite fish diversity within boundary concepts.	126
Figure 4.3.1. Geographic extent of CalCOFI bongo net data.	127

Figure 4.3.2. CalCOFI bongo net effort data.	127
Figure 4.3.3. Estimated mean larval abundance for California halibut in CalCOFI bongo tows.....	128
Figure 4.3.4. Estimated mean larval abundance for bocaccio in CalCOFI bongo tows.	129
Figure 4.3.5. Estimated mean larval abundance for Pacific sardine in CalCOFI bongo tows.....	130
Figure 4.3.6. Estimated mean larval abundance for northern anchovy in CalCOFI bongo tows.....	130
Figure 5.1.1. Ashy storm-petrel. At-sea densities (individuals/km ²) calculated for five minutes of latitude by five minutes of longitude grid cells from aerial and shipboard survey data collected from 1975-1997 and summarized in the Computer Database Analysis System v2.1 (MMS, 2001)	139
Figure 5.1.2. Ashy storm-petrel. Colony counts (corrected total individuals) 1989-1991	139
Figure 5.1.3. Brown pelican. At-sea densities (individuals/km ²) calculated for five minute of latitude by five minutes of longitude grid cells from aerial and shipboard survey data collected from 1975-1997 and summarized in the Computer Database Analysis System v2.1 (MMS, 2001)	139
Figure 5.1.4. Brown pelican. Colony counts (corrected total individuals) 1989-1991.....	139
Figure 5.1.5. Double-crested cormorant. At-sea densities (individuals/km ²) calculated for five minutes of latitude by five minutes of longitude grid cells from aerial and shipboard survey data collected from 1975-1997 and summarized in the Computer Database Analysis System v2.1 (MMS, 2001)	143
Figure 5.1.6. Double-crested cormorant. Colony counts (corrected total individuals) 1989-1991.....	143
Figure 5.1.7. Brandt's cormorant. At-sea densities (individuals/km ²) calculated for five minute of latitude by five minutes of longitude grid cells from aerial and shipboard survey data collected from 1975-1997 and summarized in the Computer Database Analysis System v2.1 (MMS, 2001)	144
Figure 5.1.8. Brandt's cormorant. Colony counts (corrected total individuals) 1989-1991	145
Figure 5.1.9. Pelagic cormorant. At-sea densities (individuals/km ²) calculated for five minutes of latitude by five minutes of longitude grid cells from aerial and shipboard survey data collected from 1975-1997 and summarized in the Computer Database Analysis System v2.....	146
Figure 5.1.10. Pelagic cormorant. Colony counts (corrected total individuals) 1989-1991	146
Figure 5.1.11. Western snowy plover. Designated critical habitat for the Pacific coast population of western snowy plover.	147
Figure 5.1.12. Black oystercatcher. Colony counts (corrected total individuals) 1989-1991	148
Figure 5.1.13. California least tern. Maximum number of breeding pairs 2001-2003	149
Figure 5.1.14. Pigeon guillemot. At-sea densities (individuals/km ²) calculated for five minutes of latitude by five minutes of longitude grid cells from aerial and shipboard survey data collected from 1975-1997 and summarized in the Computer Database Analysis System v2.1.	150
Figure 5.1.15. Pigeon guillemot. Colony counts (corrected total individuals) 1989-1991	151
Figure 5.1.16. Xantus's murrelet. At-sea densities (individuals/km ²) calculated for five minute of latitude by five minutes of longitude grid cells from aerial and shipboard survey data collected from 1975-1997 and summarized in the Computer Database Analysis System v2.1	152
Figure 5.1.17. Xantus's murrelet. Colony counts (corrected total individuals) 1989-1991	153
Figure 5.1.18. Xantus's murrelet. Anacapa Island colony counts (estimated total individuals per shoreline segment) 1991-2002	154
Figure 5.1.19. Xantus's murrelet. San Clemente Island colony counts (estimated total individuals per shoreline segment) 1991-2002	154
Figure 5.1.20. Xantus's murrelet. San Miguel Island colony counts (estimated total individuals per shoreline segment) 1991-2002	154
Figure 5.1.21. Xantus's murrelet. Santa Barbara Island colony counts (estimated total individuals per shoreline segment) 1991-2002	154
Figure 5.1.22. Xantus's murrelet. Santa Catalina Island colony counts (estimated total individuals per shoreline segment) 1991-2002	154

Figure 5.1.23. Xantus's murrelet. Santa Cruz Island colony counts (estimated total individuals per shoreline segment) 1991-2002	154
Figure 5.1.24. Xantus's murrelet. Telemetry relocations during 1995-1997 and kernel density surface based on the relocation points	155
Figure 5.1.25. Cassin's auklet. At-sea densities (individuals/km ²) calculated for five minutes of latitude by five minutes of longitude grid cells from aerial and shipboard survey data collected from 1975-1997 and summarized in the Computer Database Analysis System v2.1	155
Figure 5.1.26. Cassin's auklet. Colony counts (corrected total individuals) 1989-1991	156
Figure 5.2.1. The distribution of CDAS marine bird survey effort and sightings (left) and the total amount of effort within 5 minutes of latitude by 5 minutes of longitude grid cells (right) within the region from Point Arena to the U.S.-Mexico border	157
Figure 5.2.2. Estimated bird diversity from Point Arena, California, to the U.S.-Mexico border	158
Figure 5.2.3. Overlay of estimated marine bird diversity and CINMS boundary concepts	161
Figure 5.2.4. Regression of mean bird diversity and concept area	162
Figure 5.2.5. Regression of the area of highest mean bird diversity and concept area	162
Figure 6.1.1. Survey tracks for the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001	168
Figure 6.1.2. Survey tracks for the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 off central and southern California waters	169
Figure 6.1.3. Survey tracks for the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 off southern California waters	171
Figure 6.1.4. Survey effort (km of survey track) for the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1	172
Figure 6.1.5. Survey effort (kilometers of survey track) for the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1 off central and southern California waters	172
Figure 6.1.6. Blue whale. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 and the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1	173
Figure 6.1.7. Bottlenose dolphin. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 and the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1	174
Figure 6.1.8. Bottlenose dolphin (coastal population). Encounter rates (#/km) based on data from the Southwest Fisheries Science Center (SWFSC) aerial surveys 1990-2000	174
Figure 6.1.9. Common dolphin (unidentified). Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 and the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1, 1975-1997	175
Figure 6.1.10. Long-beaked common dolphin. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001	176
Figure 6.1.11. Short-beaked common dolphin. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001	176
Figure 6.1.12. Gray whale. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) aerial surveys conducted near San Nicolas (1992-1993) & San Clemente (1998-2003) islands and the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1, 1975-1997	178
Figure 6.1.13. Humpback whale. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 and the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1, 1975-1997	179
Figure 6.1.14. Killer whale. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 and the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1, 1975-1997	181

Figure 6.1.15. Pacific white-sided dolphin. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 and the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1, 1975-1997	181
Figure 6.1.16. Risso's dolphin. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 and the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1, 1975-1997.....	182
Figure 6.1.17. Dall's porpoise. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 and the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1, 1975-1997.....	183
Figure 6.1.18. Fin whale. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 and the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1, 1975-1997.....	184
Figure 6.1.19. Harbor porpoise. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 and the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1, 1975-1997.....	184
Figure 6.1.20. Minke whale. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 and the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1, 1975-1997.....	185
Figure 6.1.21. Northern right whale dolphin. Sightings and group size (where available) from the Southwest Fisheries Science Center (SWFSC) ship surveys 1991-2001 and the seven surveys of marine mammals compiled in the Computer Database Analysis System (CDAS) v2.1, 1975-1997	186
Figure 6.2.1. California sea lion. Maximum total individuals (greatest number of individuals of all sex and age classes counted for each beach area surveyed out of the three most recent SWFSC surveys 2001-2003) for (left to right) Anacapa, San Clemente, San Miguel, San Nicolas, Santa Barbara, Santa Catalina, Santa Cruz, and Santa Rosa islands	189
Figure 6.2.2. California sea lion. Maximum pups (greatest number of pups counted for each beach area surveyed out of the three most recent SWFSC surveys 2001-2003) for Anacapa, San Clemente, San Miguel, San Nicolas, Santa Barbara, and Gull islands	190
Figure 6.2.3. Harbor seal. Total number of individuals counted (all sex and age classes) at each surveyed site for the 2002 SWFSC census	191
Figure 6.2.4. Harbor seal. Total number of individuals counted (all sex and age classes) at each surveyed site for the 2002 CDFG census	191
Figure 6.2.5. Northern elephant seal. Maximum total individuals (greatest number of individuals of all sex and age classes counted for each beach area surveyed out of the three most recent SWFSC surveys 2001-2003) for San Clemente, San Miguel, San Nicolas, Santa Barbara, and Santa Rosa islands.....	192
Figure 6.2.6. Northern elephant seal. Maximum pups (greatest number of pups counted for each beach area surveyed out of the three most recent SWFSC surveys 2001-2003) for San Clemente, San Miguel, San Nicolas, Santa Barbara, and Santa Rosa islands.....	193
Figure 6.2.7. Southern sea otter. Counts summarized by 20 km shoreline segment for Fall 2001 and Spring 2002 (right) surveys conducted by CDFG, USGS-Biological Resources Division, and the Monterey Bay Aquarium	194
Figure 6.2.8. Southern sea otter. Expansion of sea otter range in California from 1938 to 1998.....	195
Figure 6.2.9. Southern sea otter. Three-year moving average of Spring sea otter survey counts since 1984. Reprinted with permission from the USGS Western Ecological Research Center. ..	195
Figure 7.2.1. Broad-scale distribution of the overlap of fish and invertebrate highly suitable habitat	202

Figure 7.2.2. Overlap of fish and invertebrate highly suitable habitat off southern California	203
Figure 7.2.3. Broad-scale distribution of bird and mammal co-occurrence	203
Figure 7.2.4. Bird and mammal co-occurrence off southern California	204
Figure 7.2.5. Broad-scale distribution of the composite spatial integration for bird, fish, invertebrate, and mammal data	204
Figure 7.2.6. Composite spatial integration of bird, fish, invertebrate, and mammal data off southern California.....	205

